arXiv:1108.1317v2 [astro-ph.CO] 12 Dec 2011

Cosmology with Gravitational Waves: statistical inference of the Hubble constant

Walter Del Pozzo
Nikhef, National Institute for Subatomic Physics,
Science Park 105, 1098 XG Amsterdam, The Netherlands
(Dated: today)

The precise measurement of the Hubble constant H_0 is among the foundations of the current cosmological paradigm. Due to correlations between H_0 and the remaining cosmological parameters, a precise measurement of H_0 is critical in view of future high redshift surveys. By 2015, second generation ground-based laser interferometers are expected to deliver a wealth of gravitational waves (GW) events from coalescing compact binaries up to a redshift of about 0.3. Being free of the systematics affecting electromagnetic measurements, GW offer the possibility of an independent measurement of H_0 with great accuracy. This work presents a general method based on Bayesian inference aimed at estimating the value of the cosmological parameters for any GW event. In contrast to earlier work, this framework does not require the precise identification of the putative optical counterpart, but it considers all the potential galaxy hosts consistent with the measured sky position and distance posterior distributions. When applied to the planned worldwide network of second generation interferometers, 50 GW events will yield a measurement of H_0 with an uncertainty of few percent.

PACS numbers: 95.85.Sz,98.80.-k,04.30.-w

I. INTRODUCTION

The determination of the Hubble parameter is among the most critical issues in modern cosmology, with repercussions on a wide range of fields in physics ranging from the determination of the abundances of light elements in the early Universe, the content of weakly interacting particles and therefore about the nature of dark energy. While a great deal has been learnt from the original discovery of Hubble [13], the exact value of the expansion rate today is still a matter of debate. Its current accepted value lies in the range $60-75$ km s^{-1} Mpc^{-1}[see 15, for a review] with different methods and assumptions yielding different values. For example, Cepheid variable stars in conjunction with SNe Ia observations yield $H_0 = 73 \pm 4(\text{statistical}) \pm 5(\text{systematic})$ km s^{-1} Mpc^{-1} [24], while the 15 years long Hubble Key Project [25] yields $H_0 = 62.3 \pm 1.3(\text{statistical}) \pm 5(\text{systematic})$ km s^{-1} Mpc^{-1}. Observations of the Cosmic Microwave Background from the Wilkinson Anisotropy Probe (WMAP) combined with the assumptions of a spatially flat Universe and a constant dark energy equation of state yields $H_0 = 73 \pm 3$ km s^{-1} Mpc^{-1}[17]. Without these assumptions, the data offer no additional constraints on H_0 due to the correlation with other cosmological parameters. The most precise measurement to date of the (present day) Hubble expansion rate is obtained by combining WMAP data with measurements of the Baryonic Acoustic Oscillations (BAO): $H_0 = 70.2 \pm 1.4$ km s^{-1} Mpc^{-1} [17]. Nevertheless, once one relaxes the assumption that dark energy equation of state is constant in time, H_0 is found to lie within the range 61 and 84 km s^{-1} Mpc^{-1}at the 2σ confidence level [33]. The strong correlations between H_0 and the other cosmological parameters implies that uncertainties on H_0 also affect estimates of the other parameters. All these methods yield fairly narrow statistical errors but are still affected by significant systematic uncertainties. Observations based on these established techniques are expected to significantly clarify the picture over the next few years, but a precise measurement of the Hubble parameter currently remains an open issue, and as such blurs our understanding of the high redshift universe, and its mass-energy content.

The observation of gravitational waves (GW) from coalescing compact binaries potentially offers a one-step-only, totally independent measurement of the Hubble (and other cosmological) parameters, as pointed out by Schutz [30] over 25 years ago. In GW observations the luminosity distance is a direct observable [8, 26, 30], and if one could infer from other means the redshift of the source, one could trivially estimate the cosmological parameters from the distance-redshift relation. As second-generation (or advanced) ground-based gravitational-wave laser interferometers are being installed, this becomes a very concrete scenario, which may play a decisive role in the debate. The challenge in GW observations is to obtain a redshift measurement for a GW detected source. As the GW error-box is $\approx 1-100$ deg^2, direct redshift measurements may be very challenging, despite optimistic assumptions made by several authors [*e.g.* 12, 21, 27, 38, and many others]. For example, if short-lived Gamma Ray Bursts (GRB) are associated to the merger of compact objects with at least a neutron star component, this would provide such a measurement [21, 27, 38]. However, regardless of the still open debate on whether short GRB progenitors are indeed compact binaries, the fraction of coalescing systems producing a short GRBs might be as low as 10^{-2} [3].

These considerations bear the question of whether it is at all feasible to use GW binaries as a new class of standard candles if the electro-magnetic counterpart it is not known. Recent literature focused on this possibility. Taylor et al. [32] explored the case in which the mass function of neutron stars is known, while Messenger & Read [20] suggest a direct measurement of the redshift from GWs, but the required sensitivity is achievable only by third generation instruments. The case for space-based antennas has been investigated in MacLeod & Hogan [19] for the case of H_0 and in Petiteau et al. [22] for w, the Dark Energy equation of state parameter.

In this study, I present a general method that formalises in a rigorous way the original idea proposed in Schutz [30]. The aim is to estimate the value of any cosmological parameter without assuming an electromagnetic identification of the source or a particular form of the neutron star equation of state. The results will be specialised to the case of ground-based gravitational waves detectors and I will show how they could bring a measurement of the Hubble parameter with good accuracy. The method is based on correlating a galaxy catalogue of potential hosts of the detected GW source with the error box in which one out of many is the possible host and on the combination of the results from multiple GW detections. In fact, for second-generation instruments – Advanced LIGO [16], Advanced Virgo [5] and the Large Cryogenic Gravitational-wave Telescope [18] are expected to provide several detections – the detection rate is estimated to be in the range $\sim 1-100$ yr^{-1}, depending on the actual astrophysical event rate, instrument duty cycles and sensitivity evolution [1]. The rest of the paper is organised as follows: in Section II I describe the inference model and I specialise it to the case of compact binary coalescences observed from second generation interferometers. Section III presents the GW events catalogue on which the simulation in Section IV is based. Finally, the results are summarised and discussed in Section V.

II. METHOD

Consider a catalogue of gravitational wave events $\mathcal{E} \equiv \epsilon_1, \ldots, \epsilon_n$. The posterior probability distribution for the cosmological parameters – the Hubble constant H_0, the density parameters Ω_m, Ω_k, and Ω_Λ, and so on – that I collectively represent with the $\vec{\Omega}$, for any given event ϵ_i and cosmological model $\mathcal{H}_{\vec{\Omega}}$ is given by:

$$\begin{aligned} p(\vec{\Omega}|\epsilon_i, \mathcal{H}_{\vec{\Omega}}, \mathcal{I}) &= \\ &= \int d\vec{\theta} dz \, p(\vec{\Omega}, \vec{\theta}, z|\epsilon_i, \mathcal{H}_{\vec{\Omega}}, \mathcal{I}) \end{aligned} \qquad (1)$$

where the $\vec{\theta}$ is the set of parameters on which the GW waveform depends and $\mathcal{I}$ represents all the background information relevant for the problem under scrutiny. Furthermore, the integral in (1) is to be performed over the whole parameter space. For non-spinning waveforms $\vec{\theta} \equiv (\mathcal{M}, \eta, \phi_c, t_c, \alpha, \delta, \psi, \iota, D_L)$: the *chirp* mass $\mathcal{M}$, the symmetric mass ratio η, the phase ϕ_c and time of coalescence t_c, right ascension α, declination δ, polarisation angle ι, inclination angle ι and luminosity distance D_L. Thanks to the product rule, the integrand in Eq. (1) can be factorised as:

$$\begin{aligned} & p(\vec{\Omega}, \vec{\theta}, z|\epsilon_i, \mathcal{H}_{\vec{\Omega}}, \mathcal{I}) = \\ & p(\vec{\Omega}|\vec{\theta}, z, \epsilon_i, \mathcal{H}_{\vec{\Omega}}, \mathcal{I}) p(z|\vec{\theta}, \epsilon_i, \mathcal{H}_{\vec{\Omega}}, \mathcal{I}) p(\vec{\theta}|\epsilon_i, \mathcal{H}_{\vec{\Omega}}, \mathcal{I}) \end{aligned} \qquad (2)$$

where each of the factors has a simple interpretation: (i) $p(\vec{\theta}|\epsilon_i, \mathcal{I})$, the joint probability distribution for the source parameters as inferred with parameter estimation codes [*e.g.* 34], (ii) $p(z|\vec{\theta}, \mathcal{H}_{\vec{\Omega}}, \mathcal{I})$, the probability distribution for the redshift z given the observed parameters $\vec{\theta}$ and the assumed cosmological model $\mathcal{H}_{\vec{\Omega}}$ and (iii) $p(\vec{\Omega}|\vec{\theta}, z, \epsilon_i, \mathcal{H}_{\vec{\Omega}}, \mathcal{I})$, the distribution for $\vec{\Omega}$ given the observed parameters of the source $\vec{\theta}$, the corresponding redshift distribution and $\mathcal{H}_{\vec{\Omega}}$. The inference problem is essentially solved once one specifies appropriate distributions. For the purpose of this paper, I will choose particularly simple ones to exemplify the workings of this method.

The posterior distribution $p(\vec{\theta}|\epsilon_i, \mathcal{I})$ is, in general, a complicated function of all the GW parameters and their correlations. Nevertheless, if one ignores these correlations it can be factorised in the product of the probability distributions for each single parameter. Since the redshift distribution will depend exclusively on D_L, α and δ, all the distributions for the remaining parameters will integrate to one. Therefore, all is left is $p(D_L, \alpha, \delta|\epsilon_i, \mathcal{I})$. I choose for this distribution a particularly simple form that ignores the correlations between these parameters:

$$p(D_L, \alpha, \delta|\epsilon_i, \mathcal{I}) \propto \exp\left[-\frac{(D_0 - D_L)^2}{2\sigma_D^2}\right] g(\alpha, \delta) \qquad (3)$$

where $g(\alpha, \delta)$ is given by:

$$g(\alpha, \delta) \propto \begin{cases} 1 \text{ if } \sqrt{(\delta - \delta_0)^2 + \cos^2(\delta_0)(\alpha - \alpha_0)^2} \leq \Delta\Omega \\ 0 \,\text{otherwise} \end{cases} \qquad (4)$$

where the subscript 0 indicates the mean values of the parameters of interest and $\Delta\Omega$ is the uncertainty over the solid angle. The value of $\Delta\Omega$ and σ_D can be heuristically assigned by specifying the value of the signal-to-noise ratio S/N as [9, 37]:

$$\frac{\sigma_{D_L}}{D_L} = 3 \times (S/N)^{-1} \qquad (5)$$

$$\Delta\Omega = a \times \deg^2 \left(\frac{10}{\mathrm{S/N}}\right)^2 \qquad (6)$$

where a is a factor in $\sim 10-100$ that I take to be 100. Since I neglected all the correlations among parameters and these do affect the mean value D_0 of the posterior for D_L [*e.g.* 27], I introduce a shift in D_0 of $\pm U(-1,1)\sigma_{D_L}$ with $U(-1,1)$ being a random number uniformly distributed in the interval $[-1,1]$.

Gravitational waves alone are not able to provide redshift information, therefore this piece of information needs to be acquired by other means[39]. Hence, past studies assumed a coincident GW and electromagnetic detection so to infer simultaneously distance and redshift of the source. In what follows I will relax this assumption. In this case, one can use a galaxy catalog to infer the possible values of the redshift associated with any given source. The factor $p(z|D_L, \alpha, \delta, \epsilon_i, \mathcal{H}_{\vec{\Omega}}, \mathcal{I})$, in fact, selects the regions of the z space that are consistent with the measured 3-dimensional error box and the cosmological model $\mathcal{H}_{\vec{\Omega}}$. Thus, I assign:

$$p(z|D_L, \alpha, \delta, \epsilon_i, \mathcal{H}_{\vec{\Omega}}, \mathcal{I}) = \begin{cases} \frac{1}{M}\sum_{k=0}^{M} \delta(z - z_k) \, \text{if } z_k \in [z_{\min}, z_{\max}] \text{ and } g(\alpha_k, \delta_k) \leq \Delta\Omega \\ 0 \, \text{otherwise} \end{cases} \tag{7}$$

where z_k, α_k, δ_k are the redshift and celestial coordinates of the k-th galaxy and M is the total number of galaxies in the 3-dimensional sky box. The limits $z_{\min}$ and $z_{\max}$ are determined both by the width of the distribution for D_L and by the priors for $\vec{\Omega}$. For each value of $\vec{\Omega}$ allowed by their prior distributions, the D_L–z relation predicted by $\mathcal{H}_{\vec{\Omega}}$, once evaluated at the limits of the distance distribution, gives putative values of $z_{\min}$ and $z_{\max}$. One can then choose $z_{\min}$ and $z_{\max}$ as the ones giving the largest interval.

The probability distribution $p(\vec{\Omega}|D_L, \alpha, \delta, z, \epsilon_i, \mathcal{H}_{\vec{\Omega}}, \mathcal{I})$ is instead a deterministic function. In fact, a cosmological model $\mathcal{H}_{\vec{\Omega}}$ implies that the quantities $\vec{\Omega}, D_L$ and z are not independent. For every value of the triplet $\vec{\Omega}, D_0$ and z_k there is one function f such that $f(\vec{\Omega}, D_0, z_k) = 0$. In particular, $f(\vec{\Omega}, D_0, z_k) = D_0 - D(\vec{\Omega}, z_k) = 0$. The function $D(\vec{\Omega}, z)$ is the luminosity distance expression as a function of the redshift z and of the cosmological parameters as dictated by $\mathcal{H}_{\vec{\Omega}}$. Hence, in full generality, $p(\vec{\Omega}|D_L, \alpha, \delta, z, \epsilon_i, \mathcal{H}_{\vec{\Omega}}, \mathcal{I}) = \delta(D_0 - D(\vec{\Omega}, z))$.

With these choices for the relevant probability distributions, for each GW event ϵ_i, Eq. (1) becomes:

$$p(\vec{\Omega}|\epsilon_i, \mathcal{H}_{\vec{\Omega}}, \mathcal{I}) = \frac{\sum_{k=1}^{M} \int dD_L \exp\left[-\frac{(D_0 - D(\vec{\Omega}, z_k))^2}{2\sigma_D^2}\right]}{\int d\vec{\Omega} \sum_{k=1}^{M} \int dD_L \exp\left[-\frac{(D_0 - D(\vec{\Omega}, z_k))^2}{2\sigma_D^2}\right]} \tag{8}$$

with the condition that $z_{\min} \leq z_k \leq z_{\max}$ and $g(\alpha_k, \delta_k) \leq \Delta\Omega$. This formalism is readily extended to the full catalogue of events $\mathcal{E}$:

$$p(\vec{\Omega}|\mathcal{E}, \mathcal{H}_{\vec{\Omega}}, \mathcal{I}) = \left[p(\vec{\Omega}|\mathcal{H}_{\vec{\Omega}}, \mathcal{I})\right]^{1-n} \prod_{i=1}^{n} p(\vec{\Omega}|\epsilon_i, \mathcal{H}_{\vec{\Omega}}, \mathcal{I}) \tag{9}$$

where $p(\vec{\Omega}|\mathcal{H}_{\vec{\Omega}}, \mathcal{I})$ is the prior probability distribution for $\vec{\Omega}$.

III. GRAVITATIONAL WAVE EVENTS CATALOGUE

I applied the formalism presented in the previous section to a mock catalogue of GW events observed by advanced interferometers. The galaxy catalogue used as baseline for the generation of the events is the Sloan Digital Sky Survey Data Release 7 (SDSS) [2]. The GW events catalogue consists of a set of 1,000 compact binary coalescence events distributed uniformly on the celestial sphere coverage of SDSS. Each galaxy is assigned equal probability of being the host of a GW event, therefore the redshift distribution of the GW follows closely the galaxy redshift distribution. The luminosity distance D_L is then calculated assuming a ΛCDM universe with $H_0 = 70$ km s^{-1} Mpc^{-1}, $\Omega_m = 0.3$ and $\Omega_\Lambda = 0.7$. The component masses of the binary system are chosen by drawing from a uniform distribution $m_1, m_2 \in (1.0, 15.0) M_\odot$. As SDSS covers about half of the northern sky, events that are close to the edge of the survey coverage are not considered. Sources were also restricted to have $z \leq 0.1$ (~ 460 Mpc in a ΛCDM cosmology). This redshift corresponds to the putative completeness limit of SDSS and, given the most plausible rates [1], should yield ~ 50 detections per year. The full event catalogue is then divided into 20 catalogues of 50 events each. Results will be presented in the form of an average over these 20 independent catalogues. For the calculation of the S/N, and therefore of the uncertainties over distance and sky position as described by Eq. (5), I assumed a network of three advanced interferometers; the two Advanced LIGOs in Hanford, WA, and Livingston, LA, and Advanced Virgo in Cascina, Italy. The sensitivity in each of the detectors is given by the High Power Zero Detuning configuration [28] (see Appendix A). The optimal S/N ratio in each instrument is given by:

$$\left(\frac{S}{N}\right)^2 = \int_{f_{\min}}^{f_{\mathrm{ISCO}}} df \, \frac{|h(\vec{\theta}, f)|^2}{S(f)} \tag{10}$$

where $S(f)$ is the power spectral density and $h(\vec{\theta}, f)$ is the Fourier transform of the gravitational wave template. For $h(\vec{\theta}, f)$, I use frequency domain templates belonging to the so-called TaylorF2 family [7]:

$$h(\vec{\theta}, f) = \mathcal{A} f^{-7/6} e^{i\Psi(\vec{\theta}, f)} \,, \tag{11}$$

where

$$\mathcal{A} \propto \frac{\mathcal{M}^{5/6}}{D_L} \,, \tag{12}$$

is the amplitude of the signal from a source at luminosity distance D_L, and the phase $\Psi(\vec{\theta}, f)$ at the post^2-Newtonian order is given by [4]:

$$\begin{aligned} \Psi(\vec{\theta}, f) \quad &= 2\pi f t_c - \Phi_c + \tfrac{3}{128}(\pi M f)^{-5/3}\eta^{-1}\Big[1 + \\ &+ \left(\tfrac{3715}{756} + \tfrac{55}{9}\eta\right)(\pi M f)^{2/3} - 16\pi(\pi M f) + \\ &+ \left(\tfrac{15293365}{508032} + \tfrac{27145}{504}\eta + \tfrac{3085}{72}\eta^2\right)(\pi M f)^{4/3}\Big] \end{aligned} \tag{13}$$

In the previous expressions f is the GW frequency,

$$\begin{aligned} M &= m_1 + m_2 && (14)\\ \eta &= \frac{m_1 m_2}{M^2} && (15)\\ \mathcal{M} &= \eta^{3/5} M\,; && (16) \end{aligned}$$

are the total mass, the symmetric mass ratio and the "chirp" mass, respectively, of a binary of components masses m_1 and m_2. Lastly, Φ_c is the constant phase at some (arbitrary) reference time t_c. The limits of integration in Eq. (10) are $f_{\min} = 10\text{Hz}$ and the frequency of the innermost stable circular orbit:

$$f_{\mathrm{ISCO}} = \left(6^{3/2}\pi(m_1+m_2)\right)^{-1}. \qquad (17)$$

The network S/N is then the rms of each detector S/N. Consistently with current detection searches, I considered only events yielding a S/N of 8 in each single interferometer. This results in a cut-off of 13.3 for the network S/N. Fig. 1 shows distributions for the properties of the GW catalogue. The network S/N ranges between 13.3 and ~ 60, with most of the events close to threshold. This implies that typical values for the uncertainties are $\sim 25\%$ on D_L and ~ 30 deg^2 on the sky position. These numbers are comparable with the uncertainties from second order Fisher Matrix estimates [35]. The typical number of galaxies identified as potential hosts for each GW event ranges from a *few* to $\sim 10{,}000$ depending on the S/N ratio of the source. Note that each galaxy has the same probability of hosting a GW event, so the spacial distribution of sources follows closely the spacial distribution of galaxies. Galaxies are usually found in bound structures where the typical galaxy number density is hundred times higher than the field average, therefore sources are more likely to be observed from large scale structures such as clusters or groups of galaxies having approximately the same redshift. For each event, the most probable value for $\vec{\Omega}$ will be given by the most represented redshift in the corresponding galaxy catalogue. The remaining redshifts will contribute as noise at the tails of the target posterior distribution. The clustering of galaxies was in fact the basis of the study by Schutz [30]. As in the aforementioned study, the combined PDF in Eq. (9) will effectively average out the tails of the posterior distribution for $\vec{\Omega}$, thus allowing a more accurate inference over these parameters.

IV. RESULTS

The cosmological hypothesis $\mathcal{H}_{\vec{\Omega}}$ considered for the analysis is a Friedmann-Robertson-Walker-Lemaître universe depending on four parameters, $\vec{\Omega} \equiv h, \Omega_m, \Omega_k, \Omega_\Lambda$ with the condition $\Omega_k + \Omega_m + \Omega_\Lambda = 1$. In practice, the parameters of interest are $h, \Omega_m, \Omega_\Lambda$. As customary, I defined $h \equiv H_0/100$ km s^{-1} Mpc^{-1}. The relevant expression for the luminosity distance $D(\vec{\Omega}, z)$ can be found in [11]. I choose prior probability distributions that are uniform in each parameter:

$$\begin{aligned} h &\in [0.4, 1.0] && (18)\\ \Omega_m &\in [0.0, 1.0] && (19)\\ \Omega_\Lambda &\in [0.0, 1.0]. && (20) \end{aligned}$$

For each event, I computed the posterior distribution for $\vec{\Omega}$, Eq. (8), using a Nested Sampling algorithm [29]. This algorithm evaluates directly the denominator in Eq. (8), the *evidence*, and, as a by-product, the posterior distributions of interest. I found this technique computationally more feasible that direct evaluation on a grid and, furthermore, more easily extendable to more general or complicated cosmological models. Figs. 2 and 3 show posterior distributions for $\vec{\Omega}$ for two representative events with low S/N and high S/N. In both cases, neither Ω_m or Ω_Λ is measurable, their posterior distributions are essentially flat just like their priors. For the redshift range under consideration ($z \leq 0.1$) the luminosity distance for different values of either Ω_m or Ω_Λ varies by $\sim 1\%$, which is much smaller than the typical 30% uncertainty on D_L given the S/N ratios. On the contrary, in both cases h is measurable, even though with very different accuracies. In the low S/N case, Fig. 2, the posterior for h tends to prefer values ≤ 0.7, but with a significant support at the edges of the allowed range. Note that, since most of the potential hosts have redshifts towards the upper boundary of the allowed range from the priors on $\mathcal{H}_{\vec{\Omega}}$, the probability distribution for h is skewed towards small values. A similar effect can be noted in the high S/N scenario, Fig. 3, even though since the 3-dimensional error box is very small, the posterior distribution for h is fairly well peaked around the correct value and it has very little support close to the edges of the prior range. Nevertheless, if one would estimate h from this single event, which is indeed extremely optimistic, one would infer $h = 0.7 \pm 0.1$ at the 1σ level. The combination of the information coming from multiple sources as in Eq. (9), can significantly improve our inference on h. Fig. 4 shows the evolution of the joint posterior distribution for a particular realisation of 50 GW events. The comparison between the single event posterior, in dark gray, and the joint posterior, in light gray, highlights the effect of averaging over the galaxy noise at the tails of the distribution.

h can be well determined with just few tens of observations; ~ 25 observations will yield an estimate of the Hubble parameter as $h = 0.70^{+0.04}_{-0.05}$ at 95% confidence. Combining 50 observations the estimate is improved to $h = 0.71^{+0.035}_{-0.035}$ at 95% confidence (*cfr.* Fig. 5). The results presented here qualitatively agree with the prediction by Schutz [30] that h is determined at the few percent level after $10 - 12$ observations. Nevertheless, the uncertainty on the estimate of h does not improve as the square root of the number of observations, as suggested by the aforementioned study, but it is limited by the smallest σ_{D_L}, and therefore by the largest S/N ratio, that the instruments can achieve.



FIG. 1: Properties of the GW events catalogue. *Top left*: distribution of the relative error on D_L. *Top right*: distribution of the solid angle accuracy. *Center left*: distribution of recovered luminosity distances. *Center right*: distribution of the redshift of the sources. *Bottom left*: distribution of the network S/N. *Bottom right*: distribution of the number of galaxies associated as potential hosts to each GW event.

V. CONCLUSIONS

Second generation interferometers are scheduled to start operation in 2014–2015. Within few years of observations, they are expected to deliver a wealth of GW events. GW as *standard sirens* offer the very realistic possibility of measuring H_0 in way that is independent of the "cosmic scale ladder" traditionally used in EM-based cosmology. This will be a milestone in the path of understanding the ΛCDM cosmological paradigm and our Universe in general. This paper presented a fully general method aimed at estimating the value of the cosmological parameters for any cosmological hypothesis $\mathcal{H}_{\vec{\Omega}}$, given a 3-dimensional joint probability distribution in luminosity distance D_L and sky position and a probability distribution for the redshift consistent with both the measured D_L and $\mathcal{H}_{\vec{\Omega}}$.

The application to second generation GW experiments shows that they can deliver a measurement of the Hubble parameter which is comparable with the current value derived from WMAP 7-year observations. Differently from most literature, this inference model makes use of *all* GW



FIG. 2: Posterior distributions for a sample GW event. The event S/N is 15.7 giving $D_L = (196 \pm 80)$ Mpc, $\Delta\Omega = 40$ deg^2 and a total of 1,749 potential host galaxies. The source redshift is 0.0477. *Top left*: distribution of the redshift of the potential hosts. *Top right*: posteriors for Ω_m. *Bottom left*: posteriors for h. *Bottom right*: posteriors for Ω_Λ.

events, without a need for the simultaneous observation of an EM counterpart. A scenario where one considers exclusively events with a putative EM detection neglects a significant part of the cogent information for the inference on $\vec{\Omega}$. Nevertheless, most of the information is gained from sources whose S/N ratio is high enough to provide a reasonably small error box in the sky and thus a peaked posterior distribution for $\vec{\Omega}$. Therefore, just one sure identification of the host galaxy redshift obtained, for example, by the simultaneous observation of a GRB and a GW will bring enough information to essentially drive our knowledge of $\vec{\Omega}$.

When compared with current EM measurements, GWs will not significantly improve the accuracy with which H_0 is known. The WMAP+BAO measurement yields an uncertainty of $\sim 2\%$ at 68% confidence [17]. Nonetheless, GW and EM methods are affected by *entirely different systematics*. The simulation presented in this study highlights GW as a mean of independently testing the current cosmological paradigm. Therefore, GW-based methods are to be considered not competitive, but *complementary* to EM-based ones. GW are not free from systematics. The measurement of $p(D_L, \alpha, \delta|\epsilon_i, \mathcal{I})$, for example, is known to be affected by calibration errors that, nevertheless, seem to be in general small [36]. Also the waveform family adopted for the analysis might introduce systematic effects, although, for the low mass range considered in this study, waveforms are known with sufficient accuracy to yield adequate measurements [7].

Being a proof-of-principle, this study is characterized by some simplistic prescriptions that, in a realistic case, need to be dealt with carefully. In general, the measurement of D_L cannot be disentangled from the inclination angle or from the sky position. This raises the issue of the number of detectors in operation worldwide. For instance, with only two detectors it will not be possible to measure D_L and sky position and thus to infer $\vec{\Omega}$. At the same time, if the number of detectors is greater than three, including LCGT [18] or even LIGO south [6], the accuracy for $p(D_L, \alpha, \delta|\epsilon_i, \mathcal{I})$ is greatly improved [*e.g.* 31, 35] and also the degeneracy between D_L and ι is partially broken.

The incompleteness of the survey used to obtain the redshifts will also affect the PDF for $\vec{\Omega}$. If the "true" host of the GW signal is missed, the PDF for $\vec{\Omega}$ yields a low probability for the correct values, thus contributing essentially as noise in the the joint PDF. However, unless most of the single event PDFs are biased, the joint PDF for $\vec{\Omega}$ will still converge towards their true values, with varying convergence rate. These kind of systematics are



FIG. 3: Posterior distributions for a sample GW event. The event S/N is 53 giving $D_L = (102 \pm 9)$ Mpc, $\Delta\Omega = 3.5$ deg^2 and a total of 120 potential host galaxies. The source redshift is 0.0218. *Top left*: distribution of the redshift of the potential hosts. *Top right*: posteriors for Ω_m. *Bottom left*: posteriors for h. *Bottom right*: posteriors for Ω_Λ.

especially important in preparation of the so-called third generation interferometers [23]. These instruments will observe sources in the local Universe, $z \leq 0.1$, with a S/N ratio ten times higher than advanced interferometers. Thus, they have the potential of providing an extremely accurate measurement of the PDF for H_0, possibly improving the estimate from advanced instruments by one order of magnitude. Furthermore, third generation interferometers will also observe the high redshift Universe ($z \leq 2$ for neutron star binaries) providing access to the measurement of the energy density parameters Ω_m, Ω_Λ and w. The analysis would further be complicated by the inclusion of the effect of weak lensing [10, 12]. Minimizing all the possible sources of bias in the inference of $\vec{\Omega}$ calls for extremely deep wide field sky surveys such as the planned Large Synoptic Survey Telescope (LSST) [14].

The framework presented in this study is fully general and can be applied to any GW event, regardless of the nature of the redshift information necessary for the inference based on the distance–redshift relation. This might be provided either by deep sky surveys, as has been the case for this study, or by the measurement of the tidal coupling contribution to the GW waveform [20] or by a prescription based on the neutron star mass function [32]. The inference of $\vec{\Omega}$ using gravitational waves alone, so independently of the "distance scale ladder“, offers the very concrete possibility of independently testing the current cosmological paradigm.

Acknowledgments

The author is particularly grateful to Alberto Vecchio for the invaluable discussions that led to this study. The author wish to thank T. G. F. Li, C. Van Den Broeck, S. Vitale, B. Sathyaprakash and B. Schutz for useful comments and discussions. The numerical simulations were performed on the Tsunami cluster of the University of Birmingham. This research was supported in part by the research programme of the Foundation for Fundamental Research on Matter (FOM), which is partially supported by the Netherlands Organisation for Scientific Research (NWO).

[1] Abadie, J., Abbott, B. P., Abbott, R., et al. 2010, Classical and Quantum Gravity, 27, 173001

[2] Abazajian, K. N., Adelman-McCarthy, J. K., Agüeros,



FIG. 4: Joint posterior distribution for h after combining the single event posteriors from 1, 5, 10, 15, 20, 25, 40 and 50 events, respectively, for a particular realization of 50 GW events. In dark gray the single event posterior distribution corresponding to the latest GW event included in the calculation of the joint posterior for h, in light gray. With increasing number of GW events the joint posterior becomes narrower as the noise coming from galaxies that contribute at the edges of the allowed range for h cancels out.



FIG. 5: Confidence interval evolution for H_0 as a function of the number of events considered in the analysis. The dots correspond to the posterior median value obtained from 20 realisations of 50 GW sources. The error bars correspond to the mean 95% confidence interval.

M. A., et al. 2009, ApJs, 182, 543

[3] Belczynski, K., O'Shaughnessy, R., Kalogera, V., et al. 2008, ApJL, 680, L129

[4] L. Blanchet, T. Damour, B. R. Iyer, C. M. Will and A. G. Wiseman, Phys. Rev. Lett. **74**, 3515 (1995)

[5] Bondu, F. 2010, Gravitation and Fundamental Physics in Space,

[6] Brooks, A. F., Wen, L., & Whitcomb, S. 2011, Bulletin of the American Astronomical Society, 43, #432.23

[7] Buonanno, A., Iyer, B. R., Ochsner, E., Pan, Y., & Sathyaprakash, B. S. 2009, Phys. Rev. D , 80, 084043

[8] Chernoff, D. F., & Finn, L. S. 1993, ApJL, 411, L5

[9] Cutler, C., & Flanagan, É. E. 1994, Phys. Rev. D , 49, 2658

[10] Dalal, N., Holz, D. E., Hughes, S. A., & Jain, B. 2006, Phys. Rev. D , 74, 063006

[11] Hogg, D. W. 1999, arXiv:astro-ph/9905116

[12] Holz, D. E., & Hughes, S. A. 2005, Astrophys. J. , 629, 15

[13] Hubble, E. 1929, Proceedings of the National Academy of Science, 15, 168

[14] Ivezic, Z., Tyson, J. A., Axelrod, T., et al. 2009, Bulletin of the American Astronomical Society, 41, #460.03

[15] Jackson, N. 2007, Living Reviews in Relativity, 10, 4

[16] Kissel, J. S., & LIGO Scientific Collaboration 2011, Bulletin of the American Astronomical Society, #410.07

[17] Komatsu, E., Smith, K. M., Dunkley, J., et al. 2011, ApJs, 192, 18

[18] Kuroda, K., & LCGT Collaboration 2010, Classical and Quantum Gravity, 27, 084004

[19] MacLeod, C. L., & Hogan, C. J. 2008, Phys. Rev. D , 77, 043512

[20] Messenger, C., & Read, J. 2011, arXiv:1107.5725

[21] Nissanke, S., Holz, D. E., Hughes, S. A., Dalal, N., & Sievers, J. L. 2010, Astrophys. J. , 725, 496

[22] Petiteau, A., Babak, S., & Sesana, A. 2011, Astrophys. J. , 732, 82

[23] Punturo, M., Abernathy, M., Acernese, F., et al. 2010, Classical and Quantum Gravity, 27, 194002

[24] Riess, A. G., Li, W., Stetson, P. B., et al. 2005, Bulletin of the American Astronomical Society, 37, #180.10

[25] Sandage, A., Tammann, G. A., Saha, A., et al. 2006, Astrophys. J. , 653, 843

[26] Sathyaprakash, B. S., & Schutz, B. F. 2009, Living Reviews in Relativity, 12, 2

[27] Sathyaprakash, B. S., Schutz, B. F., & Van Den Broeck, C. 2010, Classical and Quantum Gravity, 27, 215006

[28] D. Shoemaker (LSC, 2009), `https://dcc.ligo.org/cgi-bin/DocDB/ShowDocument?docid=2974`

[29] Skilling, J. 2004, American Institute of Physics Conference Series, 735, 395

[30] Schutz, B. F. 1986, Nature (London) , 323, 310

[31] Schutz, B. F. 2011, Classical and Quantum Gravity, 28, 125023

[32] Taylor, S. R., Gair, J. R., & Mandel, I. 2011, arXiv:1108.5161

[33] Tegmark, M., Strauss, M. A., Blanton, M. R., et al. 2004, Phys. Rev. D , 69, 103501

[34] Veitch, J., & Vecchio, A. 2010, Phys. Rev. D , 81, 062003

[35] Vitale, S., & Zanolin, M. 2011, arXiv:1108.2410

[36] Vitale, S., Del Pozzo, W., Li, T. G. F. et al. 2011, to be submitted to Phys. Rev. D

[37] Wen, L., & Chen, Y. 2010, Phys. Rev. D , 81, 082001

[38] Zhao, W., van den Broeck, C., Baskaran, D., & Li, T. G. F. 2011, Phys. Rev. D , 83, 023005

[39] This is not strictly true. Recently Messenger & Read [20] has shown that when at least one of the components of the binary system is a neutron star the dynamics of the tidal disruption during the late stages of the merger process in concert with the knowledge of the equation of state allows an estimate of the redshift of the source, therefore disregarding the need for auxiliary measurements. Nevertheless, the sensitivity required for a clear detection of such very high post-Newtonian effects is beyond the reach of second generation instruments with which I am concerned here.

Appendix A: Advanced LIGO Zero Detuning High Power noise curve

The noise curve used to calculate the S/N ratio is the Zero Detuning High Power design sensitivity. The analytic fit is given by:

$$\begin{aligned}\frac{S(f)}{S_0} = \quad & 60000.0\left(\frac{f}{10\mathrm{Hz}}\right)^{-29} + 5.0\left(\frac{f}{50\mathrm{Hz}}\right)^{-5} + \\ & 1.07\left(\frac{f}{100\mathrm{Hz}}\right)^{-2.25} + 3.7\left(\frac{f}{200\mathrm{Hz}}\right)^{-0.25} + \\ & 0.9\left(\frac{f}{300\mathrm{Hz}}\right)^{0.92} + 0.85\left(\frac{f}{1000\mathrm{Hz}}\right)^{1.8} + \\ & 0.35\left(\frac{f}{2000\mathrm{Hz}}\right)^{4} \qquad \text{(A1)}\end{aligned}$$

with $S_0 = 1.35 \times 10^{-50}$. The curve is shown in Fig. 6.



FIG. 6: The Advanced LIGO Zero Detuning High Power noise curve analytic fit, Eq. (A1).